

Compounded Anhydrous Sublingual Semaglutide Suspension

Ingredients: Semaglutide and Submagna™ HMW

Warning: Semaglutide possesses a risk of thyroid c-cell tumors. This drug has been shown to cause thyroid cancer in some animals, however; it is unknown if Semaglutide causes thyroid c-cell tumors in humans. Semaglutide is contraindicated in patients with a personal or family history of multiple endocrine neoplasia syndrome type 2 (MEN 2) or thyroid cancer.

Uses: Semaglutide could be used to lower blood glucose in adults with type 2 diabetes. It could also be used for promoting weight loss in adjunct with diet and increased physical activity. It also is being studied in a number of other areas including, but not limited to, the following: type-1 diabetes, overindulgence, addiction, alcoholism, memory issues, kidney disease, liver disease, lung disease, nicotine intake and others. A massive number of clinical trials are underway related to semaglutide that can be found at www.clinicaltrials.gov.

How to Take this Medication: Use this medication as prescribed. Typical administration is to place the indicated volume under the patient's tongue and be allowed to absorb there for a minimum of 90 seconds then swallow – allowing to sit under the tongue for up to 5 minutes could even be more ideal. Consider taking medication during morning/evening shower or at bedtime after completing your nighttime routine. To ensure optimal absorption and efficacy, do not eat or drink anything for 30 minutes after taking sublingual semaglutide.

Side Effects: Most commonly reported side effects when taking Semaglutide (orally) are: nausea, abdominal pain, diarrhea, decreased appetite, vomiting, and constipation. Signs of an allergic reaction present as rash, hives, itching, blistered or peeling skin, tightness in chest, swelling of lips/tongue/throat, difficulty swallowing, or difficulty breathing. If you experience any of these symptoms, please contact your doctor immediately.

Precautions: Pancreatitis has been reported in patients taking semaglutide. Discontinue and contact your doctor if you experience severe stomach pain, back pain, and/or vomiting. Any signs of visual disturbances, low blood sugar, and changes in urine should be reported to your doctor before continuation of semaglutide.

Overdose: If an overdose is identified, please contact the pharmacy or call 911. Poison control can also be reached at 800.222.1222

Missed Dose: If a dose is missed, do not try to "make up" the dose. Go on to the next scheduled dose and move forward accordingly. Do not take 2 doses at the same time or extra doses.

Storage: Store this medication in a cool, dry place. Avoid exposure to excessive heat, sunlight, or dampness.